



Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

Ramadhanil Pitopang¹⁾, Irmasari Thaha²⁾, Andi Lagaligo Amar³⁾ and Chien C. Lee⁴⁾

¹⁾Department of Biology, Faculty of Mathematics and Natural Sciences, Tadulako University, Kampus Bumi Tadulako Tondo, Jl. Soekarno Hatta Km 8, Tondo Palu, Indonesia. Tel./ Fax : +62 (0451) 422 844, Mobile : +62 852 8856 0451,

²⁾Department of Forest Management, Faculty of Forestry, Tadulako University. Kampus Bumi Tadulako Tondo, Jl. Soekarno Hatta Km 8, Tondo Palu, Indonesia

³⁾ Department of Animal Husbandry, Faculty of Animal Husbandry and Fishery Sciences, Tadulako University

⁴⁾Peti Surat P.60, Pejabat Pos Pending 93457 Pending, Sarawak, Malaysia
Tel: +60 (0) 19 8588817

ABSTRACT

Nepenthes pitopangii Chi C. Lee, Mc Pherson, Bourke & M. Mansur (Nepenthaceae), is an endemic pitcher plant to Sulawesi. The population of this plant at its type locality is only two individuals, on the other hand human activities like rattan and dammar collecting by local community and ecotourism activity have contributed to a decline in the population. The purpose of this research is to assess the current condition of the habitat of *Nepenthes pitopangii* at its type locality in Lore Lindu National Park. Field work was done in the period of September 2016-September 2017. The vegetation of *Nepenthes pitopangii* habitat was studied by using botanical exploration methods. The result showed that both location had been heavily damaged due to habitat human activities with no further evidence of the presence of *N pitopangii* at either site. Additionally, We noted a number of plant species around the habitat which were widespread in montain forest of Lore Lindu National Park such as *Xanthomyrtus angustifolia* (Myrtaceae), *Weinmania lucens* (Cunoniaceae), *Vaccinium palawanensis* (Ericaceae), *Phyllocladus hypophylus* (Podocarpaceae), *Drymis piperita* (Proteaceae), *Gleichenia vulcanica*, *Stricherus truncatus*, *Lycopodiella cernua* and *Dicksonia blumei*.

Key words: *Nepenthes pitopangii*, rare and endemic plant, Lore Lindu National Park, Central Sulawesi

INTRODUCTION

Nepenthes spp are unique and beautiful flowering plant. Locally in Malesian region, some species are utilized for cooking especially rice dishes, medicinal uses and for making rope. Many of *Nepenthes* species are cultivated and across the world as ornamental plants. Cheek and Jebb (2001) reported eight (8) species of Nepenthaceae for Sulawesi, of which four species are endemic. These numbers are comparatively low compared to the neighboring island of Borneo, where there are more than 30 species, of which 70% are endemic. Less species of *Nepenthes* in Sulawesi may be due to a lack of botanical exploration in Sulawesi, but in recent years several new species have been described from Sulawesi, such as *Nepenthes pitopangii* (Lee et al 2009), *Nepenthes undulatifolia* (Lee et al, 2009 in Mc Pherson 2009), *Nepenthes nigra* (Nerz et al, 2009 in Mc Pherson 2009) and *Nepenthes minima* (Cheek and Jebb 2016).

Nepenthes pitopangii Lee, Mc Pherson, Bourke & Mansur, endemic to the Indonesia island of Sulawesi. It was first collected by the author with Holotypus: RP. 2054 (CEB) dated 30 May 2007 from Lore Utara District, Poso Regency in the montane forest of Lore Lindu National Park, Central Sulawesi. The species is named for Dr. Ramadhanil Pitopang, a curator of the Herbarium

Celebense (CEB) Universitas Tadulako Palu, who has studied the flora of Central Sulawesi for over 18 years (Lee et al. 2009).

Nepenthes pitopangii is represents an interesting contribution to the flora of Sulawesi, together with a number of other plants which is recently described as a new species for science that comes from Sulawesi in particular of biogeography Wallacean region. On the other hand, human activities such; i) forest conversion into cacao plantation (Steffan-Dewenter et al, 2007), ii) rattan and dammar collecting, and iii) ecotourism activities like bird watching as well as mountanaring have lead to change on ecosystems and biodiversity in the region. Many studies have showed that the decline in the population of some species in this region, even tend to be scarce due to those human activities (Pitopang et al 2004; 2012a ; Gradstein et al 2007).

The purpose of this research is to study the current condition of the habitat of *Nepenthes pitopangii* at its original environment in Lore Lindu National Park, Central Sulawesi Indonesia.

II MATERIALS AND METHODS

Research Sites

Study area

The study is conducted in two locations as natural habitat of *N pitopangii*

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(*Ramadhanil Pitopang dkk*)

i.e. Helipad station at $01^{\circ} 18' 33.9''$ S and $120^{\circ} 18' 33.1''$ E with alt. 1900 m asl. and Torenali at $01^{\circ} 19' 788''$ S and 120°

$19^{\circ} 697''$ E with alt. 1700 m asl. Those research sites are located at Sedoa village, Lore Utara District Poso Regency (Fig.1)



Figure 1. Map of research site (yellow spot, track path toward to Mt. Rore Kautimbu) at the eastern part of the Lore Lindu National Park, Central Sulawesi, Indonesia.

Field sampling

Field work was made in the periode from September 2016 - September 2017. All plants species which are growing surrounding the habitat of *Nepenthes pitopangii* were recorded by using “botanical exploration methods” (van Balgooy and Tantra 1986). Voucher specimen for all plant material with flowers and fruits were collected in sets of at three (3) duplicates and deposited in Herbarium Celebense (CEB) and

Laboratory of Biodiversity Faculty of Mathematics and Natural Sciences Tadulako University Palu. Identification was done in the field, in CEB, and in the Laboratory of Biodiversity Faculty of Mathematics and Natural Sciences Tadulako University Palu

The diversity of wildlife such as species of avifauna and mammals were recorded based on the results of direct observation and interviews from the local community. Several environmental

parameters including air temperature, relative humidity, pH were measured. The relative humidity at the start and end of the exposure period were measured using a digital thermo-hygrometer (Corona^R Model: GL 99), whereas precipitation data obtained from Meteorological Geophysical Palu

Secies distribution

Phytogeographical of plant species were obtained from Flora Malesiana series (de Laubenfels, 1988 ; Sleumer 1966 ; Nooteboom 1988; Mabberley, Pannell and Sing 1995 ; de Wilde 2000, Kanis 1971 ; Kalkman 1993; Supadmo 1972 and

complemented and updated by checking the collection in the Herbarium Celebense (CEB), Palu.

RESULTS AND DISCUSSIONS

Habitat Characteristic

Based on the observation to the natural habitat of *Nepenthes pitopangii* at two research sites in the Lore Lindu National Park the habitat type was classified to high montane forest of Sulawesi (Whitten et al 1987). The geographical position and characteristic of habitat *Nepenthes pitopangii* Lee is presented in Table 1.

Table 1. The geographical position and characteristic of habitat *Nepenthes pitopangii* Lee in both research sites in Lore Lindu National Park.

| No | Parameter | Research sites | |
|----|-------------------------|---|---|
| | | Torenali | Helipad |
| 1 | Geographical position * | | |
| | Longitude (S) | 01 ⁰ 19' 788 | 01 ⁰ 18' 33.9" |
| | Latitude (E) | 120 ⁰ 19' 697" | 120 ⁰ 18' 33,1 |
| 2 | Altitude (m asl.) | 1700 | 1900 |
| 3 | Habitat description | Lightly disturbed natural montane forest, many large tree surrounding the habitat, soil surface very wet, associated with ferns. Habitat was disturbed by human activity, | Secondary vegetation, open area, small rock on the soil surface, associated with ferns, <i>Vaccinium</i> (Ericaceae), <i>Phyllocladus hypophyllus</i> . |
| 4 | Canopy cover (%) | 60 | 20 |
| 5 | Relative Humidity (%)** | 77 – 85% | 77 – 85% |
| 6 | Temperature (°C)** | 26-36 | 26-36 |
| | Maximum Manimum | 12-17 | 12-17 |

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

| | | | |
|---|----------------------|-------|-------|
| 7 | Rainfall (mm)** | 3,000 | 3,000 |
| 8 | Soil condition | | |
| | Nitrogen content (%) | 0.1 | 0.1 |
| | Phosphor (ppm) | 4.21 | 4.01 |

Notes: * = Direct measurement, ** = LLNP (2002), *** Saleh et al (2013)

Our survey of *Nepenthes pitopangii* focussed on two different research sites (Torenali and Helipad). Both of the sites were the location where this species first collected and found in Lore Lindu National Park. There are some different characteristics of the habitat. Research site 1 (Torenali) is very close to the track to Mt.Rorekautimbu. There are large tree species surrounding the habitat. While research site II (Helipad) is characterized by secondary vegetation on small rock on the soil surface. Both of research sites are situated in Montane forest of Lore Lindu National Park, Central Sulawesi, Indonesia. The location shows small-scale disturbances very close-by the path toward to Mt. Rore Kautimbu at the altitude 1650 m asl with relative humidity between 77-85%, daily temperature between 26-36° C (maximum), 12-17° C (minimum) and

annual rainfall 3,000 mm/year. The characteristic of soils are very rich in accumulation organic matter and classified into Glesols, Folic histosol, Folic cambisols (IUSS WRB, 2014) developed on granite rock on level terrain on gently sloping ridges or mid-slope terraces.

Plant diversity

We conducted a field exploration to check the presence of the species of *Nepenthes pitopangii* in their original habitat, as well as to study the diversity of plant species around the habitat using botanical exploration methods. There were forty (40) plant species recorded from both research sites. It was consisted of 27 species from research site 1 (Torenali) and 18 from research site II (Helipad) respectively. List of plant species is provided in Table 2.

Table 2. List of plants species that growing surround the habitat of *Nepenthes pitopangii* Lee in the studied area :

| No | Botanical Name | Family | Habitus | Sites | | Distribution |
|----|--|-------------|---------|-------|----|--------------------|
| | | | | I | II | |
| 1 | <i>Castanopsis accuminatisima</i> Rehder | Fagaceae | Tree | √ | √ | Malesia |
| 2 | <i>Lithocarpus celebicus</i> (Miq.) Rehder | Fagaceae | Tree | √ | | Sulawesi, Filipina |
| 3 | <i>Pandanus sarasinorum</i> Warb | Pandanaceae | Tree | √ | √ | Endemic Sulawesi |
| 4 | <i>Neolitsea javanica</i> (Blume) Backer | Lauraceae | Tree | √ | | West Malesia |

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

| | | | | | | |
|----|---|-----------------|------------------|---|---|--|
| 5 | <i>Adinandra celebica</i> Koord. | Theaceae | Tree | √ | | Endemic Sulawesi |
| 6 | <i>Eucalyptus deglupta</i> Blume | Myrtaceae | Tree | √ | | Sulawesi, Mindanao, Seram, papua |
| 7 | <i>Quintinia apoensis</i> (Elmer) Schltr. | Parachrypiaceae | Tree | | | Central Sulawesi, Papua |
| 8 | <i>Gastonia serratifolia</i> (Miq) Phillipson | Araliaceae | Tree | √ | √ | Malesia |
| 9 | <i>Acronychia trifoliata</i> Zoll.& Moritzi | Rutaceae | Tree | √ | | Malesia |
| 10 | <i>Pinanga caesia</i> Blume | Arecaceae | Tree | √ | | Endemic |
| 11 | <i>Acmena acuminatissima</i> (Blume) Merr. & L.M. Perry | Myrtaceae | Treelet | √ | √ | Malesia |
| 12 | <i>Litsea ferruginea</i> Blume | Lauraceae | Tree | | | West Malesia |
| 13 | <i>Drimys piperita</i> Hook.f | Winteraceae | Treelet | √ | | Sulawesi, Borneo, Papua, Malesia |
| 14 | <i>Santiria sp</i> | Burseraceae | Treelet | | | |
| 15 | <i>Weinmannia lucen</i> Baker | Cunoniaceae | Treelet | √ | | Sulawesi, Filipina |
| 16 | <i>Pittosporus mollucanum</i> (Lam) Miq. | Pittosporaceae | Treelet | √ | | |
| 17 | <i>Eurya acuminata</i> DC | Theaceae | Treelet | √ | | Malesia |
| 18 | <i>Phyllocladus hypophyllus</i> Hook | Phyllocladaceae | Tree | √ | | Sulawesi, Wallacea, Papua, New Guinea, Filipina, Kalimantan |
| 19 | <i>Dacrycarpus imbricatus</i> (Blume) de Laub | Podocarpaceae | Treelet | √ | | Malesia |
| 20 | <i>Vaccinium palawanense</i> Merr | Ericaceae | Shrub | | √ | Sulawesi, Filipina |
| 21 | <i>Psychotria celebica</i> Miq. | Rubiaceae | Shrub | √ | | Endemic Sulawesi |
| 22 | <i>Xanthomyrtus angustifolia</i> A.J. Scot | Myrtaceae | Shrub | √ | √ | Sulawesi, Papua |
| 23 | <i>Litsea ferruginea</i> Blume | Lauraceae | Treelet | | | East Malesia |
| 24 | <i>Viburnum amplificatum</i> J Kem | Caprifoliaceae | Treelet | | √ | Borneo, Sulawesi |
| 25 | <i>Macaranga cf.costulata</i> Pax & K.Hoffm | Euphorbiaceae | Treelet | √ | | |
| 26 | <i>Themeda arguens</i> L | Poaceae | Herb | | √ | |
| 27 | | Rubiaceae | Herb | | √ | |
| 28 | <i>Anaphalis javanica</i> (DC.) Sch.Bip. | Asteraceae | Herb | | √ | Sumatra, Java, Borneo, Sulawesi, PNG |
| 29 | <i>Mischocarpus sundaicus</i> Blume | Sapindaceae | Shrub | | √ | |
| 30 | <i>Agathis celebica</i> (Koord.) warb. | Araucariaceae | Tree | | √ | Sulawesi, Filipina |
| 31 | <i>Trachymene acerifolia</i> C. Norman | Umbeliferae | Herb | √ | | Lesser Sunda, Sulawesi |
| 32 | <i>Nepenthes mirabilis</i> (Lour) Rafarin | Nepenthaceae | Climber | √ | | Malesia, North Australia, Cambodia, Laos, Myanmar, China (Hainan) |
| 33 | <i>Dissochaeta celebica</i> Blume | Melastomataceae | Climber | | √ | Endemic Sulawesi |
| 34 | <i>Dipteris conjugata</i> Reinw. | | Terrestrial fern | √ | √ | Indochina, southern China, Taiwan, Malesia to New Calodonia and Australia. |
| 35 | <i>Gleichenia vulcanica</i> Blume | Gleicheniaceae | Terrestrial | √ | √ | Pantropical |

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

| | | | | | | |
|-----------|--|---------------------|------------------|----------|----------|---|
| | | | fern | | | |
| 36 | <i>Sticherus truncatus</i> (Willd.) Nakai | Gleicheniaceae | Terrestrial fern | √ | √ | Indochina, Malesia, North Australia |
| 37 | <i>Lycopodiella cernua</i> (L.) Pic. Serm | Lycopodiaceae | Terrestrial fern | √ | √ | Neotropics, Malesia |
| 38 | <i>Dicksonia blumei</i> (Kunze) Moore | Dicksoniaceae | Tree fern | | √ | Malaysia, Indonesia, Philippines and PNG |
| 39 | <i>Racemosbamboo celebica</i> Dransfield | Poaceae | Climber | √ | | Sulawesi |
| 40 | <i>Rubus mollucanus</i> L | Rosaceae | Climber | √ | | All over Malesia, Thailand, Vietnam, Fiji, Caledonia, |
| 41 | <i>Nepenthes pitopangii</i> | Nepenthaceae | Climber | X | X | Central Sulawesi |

Data source : Primary data (2017). √ = present, X = absent

The result of observation showed that there was no more *Nepenthes pitopangii* at both habitat (Table 2 and Figure 2). We noted that habitat destruction of species might be due to the fact that both sites are easily accessed by the people and visitors of the Park. Both of the habitats are located on the hiking trail to Mount Rorekautimbu. It was previously used by vehicles transporting timber by Forest Concession Company (“HPH Kebun Sari”) and helipad formerly used for helicopter landing. Both locations indicated that the habitat was disturbed by the people who crossed this place for several purposes like hiking as well as damar and rattan collection. The extinction of *Nepenthes pitopangii* might also be caused by the people who illegally collected the specimen without permission from park authority. However, it was suspected that such species still exist, but it still needs exploration effort intensively to ensure the

presence of species. The limited of time, the difficulty of terrain and the presence of military operations in the region were also a limited factor in revealing the existence of the species.

In Lore Lindu National Park, Central Sulawesi, especially in the mountain forest around the route to Mount Rorekautimbu and surrounding areas was reported as many as 5 species of *Nepenthes* namely: *Nepenthes pitopangii*, *Nepenthes tentaculata*, *Nepenthes maxima* and *Nepenthes nigra* (Lore Lindu National Park Office, 2016).

As one of the main tourist destination in Lore Lindu National Park, the area around the Rorekautimbu mountain was visited by many domestic and international tourists, because the location is very easy access from Palu, the Capital city of Central Sulawesi Province. Besides, the region is the home of the charming of Wallacean biological

diversity, especially the mountain flora and fauna of Sulawesi. (LLNP, 2016). The area was also very important for local people (ethnically called "Topo Baria") who lived around the village of Sedoa, where most people had livelihoods to collect resin of damar and rattan (Alberta M, village leader, 2018, pers. comm.)

Based on IUCN Red list (<http://www.iucnredlist.org/details/49000915/0>, 2017) and Lee et al. (2009) that conservation status of *Nepenthes pitopangii* is categorized as vulnerable under criterion D2 which lies within a protected area (PA). However, the PA site supports just one plant, which appears to be under threat from plant collectors. The area of occupancy and extent of occurrence are both 8 km². Fortunately, the other known locality supports a few hundred plants (population size is about 400 mature individuals) and it is probable that other, undiscovered subpopulations exist. Nevertheless, based on current knowledge, the conservation status of *N. pitopangii* has the potential to change suddenly given the threat of collection and the species would benefit from a systematic monitoring programme. The species is listed on CITES Appendix II.

The major vegetation of habitat *Nepenthes pitopangii* were composed by *Castanopsis acuminatisima* (Fagaceae), *Lithocarpus celebicus* (Fagaceae),

Pandanus sarasinorum (Pandanaceae), *Neolitsea celebica* (Lauraceae), *Agathis celebica* (Podocarpaceae), *Adinandra celebica* (Theaceae), *Eucalyptus deglupta* (Myrtaceae), *Phyllocladus hypophyllus*, *Quintinia apuensis*, *Gastonia serratifolia*, *Pinanga caesea* (Arecaceae), *Santiria* sp, *Acmena acuminatisima* (Myrtaceae) and *Listea feruginea* (Lauraceae). Sapling and shrub species were *Pittosporum moluccanum*, *Eurya acuminata*, *Gastonia seratifolia*, *Drymis piperita*, *Phyllocladus hypophyllus*, *Litsea feruginea*, *Weinmania* sp, *Magnolia vriescana*, *Vaccinium palawanensis* and *Xanthomyrtus angustifolius*. There were some species of herbs such as *Themeda arguens*, *Anaphalis javanica*, *Mischocarpus sundaicus* and *Trachymene acerifolia*. We also noted a pitcher plant *Nepenthes maxima*, an endemic climber bamboo (*Racemosbamboo celebica*). Some species of terrestrial fern such as *Dipteris conjugata*, *Gleichenia vulcanica*, *Stricherus truncatus*, *Lycopodiella cernua* and *Dicksonia blumei* were found.

Vegetation of habitat is the main constituent of mountain forests of Sulawesi in Lore Lindu National Park which is rich in Fagaceae-Myrtaceae, some of which are endemic to Sulawesi such as *Pandanus sarasinorum* (Pandanaceae), *Adinandra celebica* (Theaceae), *Quintinia apoensis* (Parachrypiaceae), *Pinanga caesea* dan

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(*Ramadhanil Pitopang dkk*)

Racemosbamboo celebica. Some species such as *Eucalyptus deglupta* (Myrtaceae), *Drymis piperita* (Proteaceae), *Weinmania lucen*, *Phyllocladus hypophyllus*, *Vaccinium palawanensis*, *Xanthomyrtus angustifolia* (Myrtaceae), *Litsea ferruginea* (Lauraceae), *Agathis celebica* (Araucariaceae) have limited distribution in Sulawesi, the Philippines and eastern part of Malesia. The flora of Mt Rorekautimbu was formerly known from the floristic studies of van Balgooy and Tantra (1986), Culmsee et al. (2011), Pitopang (2012^b) and Brambach et al (2016). The forest is rich in chesnut family (Fagaceae), conifers and Myrtaceae..

Phytogeography of Sulawesi Island is located in the Wallacea region, an area that has an unique biodiversity. Isolation geography of the Sulawesi island and environmental conditions, such as variations in topography, gradient elevation, and variations in soil types has allowed spesifically the development of flora and fauna in the bioregion (Siebert 2002; Pitopang 2012^b). Total species richness and endemism of Sulawesi are comparable to those of Sumatra, Java, Borneo and New Guinea, in spite of the very different geological history of Sulawesi and the greater distance of the island to the mainland (Roos et al. 2004).

Whereas the islands of Borneo, Sumatra and Java had terrestrial connections to the mainland Asia in the past, Sulawesi was always isolated from these islands as well as from New Guinea by deep maritime straits as shown by Hall (1995) through the reconstruction of the Malay archipelago since 50 million ago. Approximately 15% of the known flowering plant species of Sulawesi are endemic (Whitten et al, 1987). Van Balgooy et al. (1996) recognized 933 indigenous plant species on Sulawesi and of these 112 were endemic to the island.

Neighborhood habitat of *Nepenthes pitopangii* also rich in various kinds of wildlife from different taxa. Based on the data recorded, several types of mammals such as Highland anoa (*Bubbalus quarlesii*), Cuscus bear (*Aliurop ursinus*) and a number species of Avifauna with their natural distribution limited on the island of Sulawesi such as *Dicaeum aureolimbatum*, *Ptilinopus fischeri*, *Dicaeum nehrikorni*, *Dicrurus Montanus*, *Myza celebensis*, *Myza sarasinorum*, *Cyornis hoevelli*, *Ficedula bonthaina*, *Loriculus exilis*, *Rhapidura teysmanii*, *Scissirostrum dubium*, *Enodes erythrophris*, *Scolopax celebensis*, *Geomalia hendrichia*, *Artamus monachus* are found (Mallo 2015)



Fig. 2. Habitat of *Nepenthes pitopangii* at two research sites in the Lore Lindu National Park, Central Sulawesi. 1 (above left). Location I, Torenali, Track to Mt. Rorekautimbu, 2. The second collection of *Nepenthes pitopangii* (taken 2008), 3. There is no more *Nepenthes pitopangii* at Torenali (surveyed by author 2016), 4 (above right) Location II, Helipad. 5. *Nepenthes pitopangii*, Photograph by author (2013) at Helipad, 6. Current condition of habitat at Helipad (Surveyed by author (2017)

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

It was concluded the habitat had been heavily damaged due to habitat human activities with no further evidence of the presence of *N pitopangii* at either site. Additionally, We noted a number of plant species around the habitat which were widespread in mountain forest of Lore Lindu National Park such as *Xanthomyrtus angustifolia* (Myrtaceae), *Weinmania lucens* (Cunoniaceae), *Vaccinium palawanensis* (Ericaceae), *Phyllocladus hypophylus* (Podocarpaceae), *Drymis piperita* (Proteaceae), *Gleichenia vulcanica*, *Stricherus truncatus*, *Lycopodiella cernua* and *Dicksonia blumei*. We noted a various kinds of wildlife from different taxa in the habitat.

ACKNOWLEDGEMENTS

We would like to express our gratitude to The Ministry of Research and Technology Directorate General of Higher Education Republic of Indonesia for financial support through “Fundamental Research Program”. We also thank Donny Mangitung Ph.D and staffs who has organize this research project and the Lore Lindu National Park Authority (LLNP) for permission to conduct this research. We thank Sahlan, Dian M.Fauzan, Fajri Ramadhan and Kurniawan B.Banjolu for their help in this work.

REFERENCES

- Brambach F, Coode MJE, Biagioni S, Culmsee H. 2016. *Elaeocarpus firdausii* (Elaeocarpaceae), a new species from tropical mountain forests of Sulawesi. *PhytoKeys* 62: 1-14
- Cheek M, Jebb.MHP. 2001. Nepenthaceae. In Nooteboom HP (ed.). *Flora Malesiana* 15. 1-157. Nationaal Herbarium of Nederland, Leiden.
- Cheek, M and MHP, Jebb. 2016. *Nepenthes minima* (Nepenthaceae), a new pyrophytic grassland species from Sulawesi Indonesia. *Blumea* 61.2016. 181-186
- Culmsee H, Pitopang R, Mangopo H, Sabir S. 2011. Tree diversity and phytogeographical patterns of tropical high mountain rain forests in Central Sulawesi, Indonesia. *Biodivers Conserv J*. 20:1103–1123
- de Laubenfels DJ. Coniferales. 1988. Series 1. Vol. 10; 337-453. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- de Wilde WJJO. 2000. Myristicaceae. Series I. Vol. 14; 1-632. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Gradstein SR, Kessler M and Pitopang R. 2007. Tree species diversity relatives to human land uses in tropical rain forest margins in Central Sulawesi. . In: Tschardtke T, Leuschner C, Zeller M, Guhardja E, Bidin A (eds) *The stability of tropical rainforest margins, linking ecological, economic and social constraints of land use and conservation.*

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

- Springer, Berlin Heidelberg New York. Page; 319-332
- Hall R. 1995. The plate tectonics of Cenozoic SE Asia and the distribution of land and sea. Pages 99- 131 in Biogeography and geological evolution of SE Asia (R. Hall, and J. D. Holloway, eds.). Backhuys, Leiden
- <http://www.iucnredlist.org/details/49000915/0>. 2017. The IUCN Redlist of Threatened species. *Nepenthes pitopangii*
- IUSS Working Group WRB (2014) World Reference Base for Soil Resources 2014. International soil classification system for naming soils and creating legends for soil maps. FAO, Rome, 181 pp. <http://www.fao.org/3/a-i3794e.pdf>
- Kanis A. 1971. Ochnaceae. Flora Malesiana Series 7. Vol.1.; 97-119. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Kalkman C. 1993. Rosaceae. Flora Malesiana Series 1. Vol.11 (2); 227-351. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Keßler PJA, Bos MM, Sierra Daza SEC, Kop A, Willemse LPM, Pitopang R, Gradstein SR, 2002. Checklist of woody plants of Sulawesi, Indonesia. Blumea Suppl 14:1–160
- Kessler M, Keßler PJA, Gradstein SR, Bach K, Schnull M and Pitopang R . 2005. Tree diversity in different land use systems in Central Sulawesi, Indonesia. Biodiversity and Conservation. Lee, C.C., McPherson, S., Bourke, G. and Mansur, M. 2009. *Nepenthes pitopangii* (Nepenthaceae), a new species from central Sulawesi, Indonesia. *The Gardens' Bulletin Singapore* 61(1): 95-100.
- Lee CC, A Wistuba, J Nerz, U Zimmermann, AP Paserang and R Pitopang. 2009. *Nepenthes undulatifolia*, a new pitcher plant from South East Sulawesi Indonesia. In McPherson S.R., 2009, *Pitcher Plants of the Old World* (Volumes 1 & 2), Redfern Natural History Productions Ltd: Poole, England. 493-505
- Lore Lindu National Park Office. 2016. Statistics of Lore Lindu National Park. Taman Nasional Lore Lindu Palu. Mabberley DJ, Pannell CM and Sing AM. 1995. Meliaceae. Series 1, vol 12. (1). 1-407. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Mallo FN, 2015. Burung-burung di Taman Nasional Lore Lindu, Catatan Ekologi, Konservasi dan Status Keberadaan Jenis. Dipublikasikan oleh CBC. Diterbitkan oleh Program Studi Magister Ilmu Lingkungan (PSMIL) Universitas Padjadjaran.
- Nerz J, A Wistuba, CC Lee, G Bourke, U Zimmermanns and S Mc Person. 2009. *Nepenthes nigra*, a new pitcher plant from Central Sulawesi. In McPherson S.R., 2009, *Pitcher Plants of the Old World* (Volumes 1 & 2), Redfern Natural History Productions Ltd: Poole, England. 469-491
- Nooteboom HP. 1988. Magnoliaceae. Seri 1. vol. 10 ; 561-604. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Pitopang R, Keßler PJA, Gradstein SR, Guhardja E, Tjitrosudirdjo SS and Wiriadinata H. 2004. Tree Composition in Secondary forest of Lore Lindu National Park,

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)

- Central Sulawesi Indonesia. In: Gerold G, Fremery M, Guhardja E (eds) Land Use, Nature Conservation and the Stability of Rainforest Margins in Southeast Asia. Springer, Berlin Heidelberg New York, Page : 269-279
- Pitopang R. 2012a. Impact of forest disturbance on the structure and composition of vegetation in tropical rainforest of Central Sulawesi, Indonesia. *Biodiversitas* 13 (4) ; 179-189
- Pitopang R. 2012b. Struktur and Komposisi Vegetasi Pada 3 Zona Elevasi yang berbeda di Taman Nasional Lore Lindu, Sulawesi Tengah Indonesia. *J.Natural Science*. Vol.1 (1); 85-105
- Roos M, Keßler PJA, Gradstein SR and Baas P. 2004. Species diversity and endemism of 5 major Malesian islands: diversity-area relationships. *Biogeogr.* 31: 1893-1908
- Saleh MFR, Miswan and R Pitopang. 2013. Autekologi *Nepenthes pitopangii* di Kawasan Taman Nasional Lore Lindu Sulawesi Tengah. Vol. 2 (2): 8-18
- Siebert SF. 2002. From Shade to Sun grown Perennial Crops in Sulawesi Indonesia. Implications for Biodiversity Conservation and Soil Fertility. *Biodiversity and Conservation*. 11. 1889-1902.
- Sleumer H. 1966. Ericaceae. *Flora Malesiana Serie I. Vol. 6* ; 469-914. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Stefan-Dewenter I, M Kessler, J Barkmann, MM Bos, D Buchori, S Erasmi, H Faust, G Gerold, K Glenk, SR Gradstein, E Guhardja, M Harteveld, D Hertel, P Hohn, M Kappas, S Kohler, CH Leuschner, M Maertens, R Margggraf, S Migge-Kleian, J Mogeia, R Pitopang, M Schaefer, S Schwarse, S G Sporn, A Streingrebe, SS Tjitrosoedirdjo, S Tjitrosoemito, A Twele, R Weber, L Woltmann, M Zeller and T Tschardtke. 2007. Tradeoffs between income, biodiversity, and ecosystem functioning during tropical rainforest conversion and agroforestry intensification. *PNAS* published online Mar 14, 2007
- Supadmo E. 1972. Fagaceae. *Flora Malesiana* , Series I.vol 7.: 266-403. Published by Foundation of Flora Malesiana. Rijks Herbarium/Hortus Botanicus. Leiden University. The Netherland
- Van Balgooy MMJ, Tantra IGM. 1986. The vegetation in two areas in Sulawesi, Indonesia. *Forest Research and Development Center Bulletin*. Bogor
- Van Balgooy MMJ, Hovenkamp PH, Welzen PC. 1996. Phytogeography of the Pasific-Floristic and historical distribution pattern in plant. In *The origin and evolution of Pasific island biotas. New Guinea to eastern Polynesia ; pattern and process*. Pp. 191-213. edited by Keast A, Miller SA. SPB academic Publishing bv. Amsterda
- Whitten AJ, Mustafa M and Henderson GS. 1987. *The Ecology of Sulawesi*. Gadjah Mada University Press. Yogyakarta

Has *Nepenthes pitopangii* Chi C. Lee, S. Mc Pherson, Bourke & M. Mansur a rare and endemic flowering plant of Sulawesi, been extinct in its original nature?

(Ramadhanil Pitopang dkk)